CORRECTION Open Access

Correction: Olfactory swab sampling optimization for α-synuclein aggregate detection in patients with Parkinson's disease



Matilde Bongianni^{1†}, Mauro Catalan^{2†}, Daniela Perra^{1†}, Elena Fontana¹, Francesco Janes³, Claudio Bertolotti², Luca Sacchetto⁴, Stefano Capaldi⁵, Matteo Tagliapietra¹, Paola Polverino², Valentina Tommasini², Giulia Bellavita², Elham Ataie Kachoie⁵, Roberto Baruca⁶, Andrea Bernardini³, Mariarosaria Valente³, Michele Fiorini¹, Erika Bronzato¹, Stefano Tamburin¹, Laura Bertolasi¹, Lorenzo Brozzetti¹, Maria Paola Cecchini¹, Gianluigi Gigli³, Salvatore Monaco¹, Paolo Manganotti² and Gianluigi Zanusso^{1*}

Correction: Translational Neurodegeneration (2022) 11:37 https://doi.org/10.1186/s40035-022-00311-3

In the original publication of this article [1], "AN" is missing in the column heading "Patients underwent NS at the (n=46)" in Table 1 due to a typesetting error. The correct column heading should be "Patients underwent NS at the AN (n=46)". Moreover, the asterisk symbol * in "27 (84)" and "5 (45)" should be removed.

In addition, the phrase "using FLOQBrushes (Copan Italia, Brescia, Italy)" needs be added in the first sentence under the header "OM sample collection" of the Methods section.

The first sentence should be changed to:

NS was performed by otolaryngologists in each unit, independently, using FLOQBrushes (Copan Italia, Brescia, Italy).

The original article [1] was updated.

Author details

¹Department of Neurosciences, Biomedicine, and Movement Sciences, Policlinico G. B. Rossi, University of Verona, 37134 Verona, Italy. ²Neurology Unit, Department of Medicine, Surgery and Health Sciences, Ospedale Cattinara, University of Trieste, 34128 Trieste, Italy. ³Neurology Unit, University of Udine Academic Hospital, 33100 Udine, Italy. ⁴Department of Surgical Sciences, Dentistry, Gynecology and Pediatrics, University of Verona, 37134 Verona, Italy. ⁵Biocrystallography Laboratory, Department of Biotechnology, University of Verona, 37134 Verona, Italy. ⁶Otolaryngology Unit, Department of Medicine, Surgery and Health Sciences, Ospedale Cattinara, University of Trieste, 34128 Trieste, Italy.

Published online: 12 August 2022

Reference

 Bongianni, et al. Olfactory swab sampling optimization for α-synuclein aggregate detection in patients with Parkinson's disease. Transl Neurodegener. 2022;11:37.

The original article can be found online at https://doi.org/10.1186/s40035-022-00311-3.

†Matilde Bongianni, Mauro Catalan and Daniela Perra contributed equally to

*Correspondence: gianluigi.zanusso@univr.it

¹ Department of Neurosciences, Biomedicine, and Movement Sciences, Policlinico G. B. Rossi, University of Verona, 37134 Verona, Italy Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.